

WEBINAIRE TRACES

TRANSFORMER LA MODÉLISATION DU CLIMAT
POUR LES SERVICES CLIMATIQUES

"LES CHANGEMENTS DES SITUATIONS
MÉTÉOROLOGIQUES RENFORCENT LES IMPACTS DES
VAGUES DE CHALEURS ET TEMPÊTES EN
FRANCE ET EN EUROPE"

DAVIDE FARANDA, AGLAÉ JEZEQUEL, MATHIEU VRAC ET PASCAL YIOU













MOTIVATION Impactful extreme events are largely driven by atmospheric dynamics

03 June 2022 Heatwave over France... A recurrent situation in recent years => Role of climate change?








BOTTLENECK Detecting robust changes in atmospheric circulation is difficult


- 1. Previous studies are not conclusive on whether anthropogenic affects atmospheric dynamics changing structure, frequency and persistence of patterns.
- 2. The diversity in the results comes from the variety of indicators of atmospheric dynamics: indices or projections that could hide robust changes.
- 3. Here we **employ a technique based on the study of analogues** of weather patterns, preserving all information contained in the weather fields to **detect robust changes**.






METHODS Using analogs frequency changes to detect long-term dynamical changes

- 1.We select alternatively all ERA5
 Sea-level Pressure maps over the
 North Atlantic from 1950-2021
- 2. For each map we compute the Euclidean distance between daily maps, and define a high quantile to select the analogues.
- 3. We count the number of analogues per decade and we select those having an increasing trend and a decreasing trend









METHODS Ensuring Robustness of Changes

- 1. For each SLP map (a) we take the best 2% (b), 1% (c) and 0.5% (d) analogues in daily sea-level pressure maps from ERA5 starting in January 1950.
- 2. The **# Analogues counted in each decade** are then obtained (blue dots, panels b--d).
- 3. A linear fit is performed (solid line) and the confidence intervals for the slope are computed (dotted lines).
- 4. Only if all confidence intervals have a positive slope, the map has increasing frequency









RESULTS: WINTER


The vast majority (92.7%) of circulation patterns show no significant occurrence trend in the historical period; 5.1% show increasing trends and 2.2% show decreasing trends


INCREASING TRENDS





DECREASING TRENDS







RESULTS: WINTER


The vast majority (92.7%) of circulation patterns show no significant occurrence trend in the historical period; 5.1% show increasing trends and 2.2% show decreasing trends

INCREASING TRENDS DECREASING TRENDS 2-meter Temperature [°C] 2-meter Temperature [°C] Precipitation Rate [mm/day] Precipitation Rate [mm/day]



RESULTS: SUMMER


The vast majority (92.7%) of circulation patterns show no significant occurrence trend in the historical period; 5.1% show increasing trends and 2.2% show decreasing trends





RESULTS: SUMMER


The vast majority (92.7%) of circulation patterns show no significant occurrence trend in the historical period; 5.1% show increasing trends and 2.2% show decreasing trends








RESULTS: Role of Interannual Variability

We find a non-significant influence of AMO in the trends















RESULTS: Role of Interannual Variability

We find a moderate role of ENSO, but it is not sufficient to explain all the trends observed













RESULTS: Linking frequency changes to impactful extreme events







SUMMARY



We find that large scale atmospheric patterns which favor summertime heatwaves and wintertime windstorms over large parts of the continent are becoming increasingly frequent



We have also shown that ENSO and AMO play a minor role in these changes. These trends can be therefore linked to anthropogenic climate change.



A key implication of our work is that anthropogenically-induced circulation changes modulate extreme events already in the present climate.





JOUEZ À

CLIMA RISQ



COMPRENDRE LA COMPLEXITÉ DU SYSTÈME CLIMATIQUE ET L'URGENCE D'UNE ACTION COLLECTIVE















The analyses described here are part of two European projects EDIPI and XAIDA received funding from the European Union's Horizon 2020 research and innovation program, grants no. 956396 and no. 101003469.

Twitter: @DaviFaranda

Email: davide.faranda@cea.fr

www.davide-faranda.com



D Faranda, G Messori, A Jézéquel, M Vrac, P Yiou. Atmospheric circulation compounds anthropogenic warming and extreme climate impacts in Europe. PNAS, 2023 https://doi.org/10.1073/pnas.2214525120

















